

PLACE

ACADEMY

I Seminari di PLACE Academy 2023
Corsi ECM residenziali



Direzione scientifica
Leonardo Calò - Marco Rebecchi
Policlinico Casilino - Roma

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Dott.ssa Marina Comisso

Cardiochirurgia e trapianto di Cuore
Ospedale San Camillo-Forlanini
Roma

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Scompenso cardiaco avanzato



Trapianto di Cuore: gold standard per il trattamento dello scompenso cardiaco avanzato che non risponde a terapia convenzionale.

Pazienti già in trattamento ottimale con terapie convenzionali per l'insufficienza cardiaca


(Terapia medica ottimale, ICD, CRT, chirurgia convenzionale)

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa



Sopravvivenza Trapianto di Cuore in Italia

Casistica Adulti 2000-18											
1 mese		3 mesi		6 mesi		3 anni		5 anni		10 anni	
Paziente	Organo	Paziente	Organo	Paziente	Organo	Paziente	Organo	Paziente	Organo	Paziente	Organo
89,4%	88,8%	85,4%	84,9%	83,6%	83,1%	76,9%	76,3%	72,9%	72,2%	61,1%	60,3%
[88,5% - 90,2%]	[87,9% - 89,6%]	[84,4% - 86,3%]	[83,9% - 85,9%]	[82,5% - 84,6%]	[82,1% - 84,1%]	[75,7% - 78%]	[75% - 77,4%]	[71,6% - 74,2%]	[70,9% - 73,5%]	[59,5% - 62,6%]	[58,7% - 61,8%]




5 anni		10 anni	
Paziente	Organo	Paziente	Organo
72,9%	72,2%	61,1%	60,3%
[71,6% - 74,2%]	[70,9% - 73,5%]	[59,5% - 62,6%]	[58,7% - 61,8%]

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Sopravvivenza Trapianto di Cuore età Pediatrica e Adulti (ISHLT)

(Transplants: January 1992 – June 2017)



I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Trapianto Cardiaco in Italia




TRAPIANTI DI CUORE



Periodo 2013-2022

● Variazione % rispetto al 2021



ATTIVITÀ PER REGIONE Anno 2022

Lombardia 50

Friuli Venezia Giulia 21

Toscana 14

Basilicata 0

Veneto 42

Piemonte 19

Sicilia 12

Calabria 0

Emilia Romagna 29

Lazio 18

Sardegna 7


Liguria 0

Puglia 26

Campania 16

Abruzzo 0

Marche 0



Fonte: 2013-2021 SIT - 2022 CNT OPERATIVO (DATI PRELIMINARI)

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Lista per trapianto di cuore in Italia

Al 31 dicembre 2021, **1041 iscrizioni**; il tasso di **mortalità** in lista è pari al **6,9%** mentre l'indice di **soddisfamento di lista totale** è di **24,2%**



TMA IN LISTA DI ATTESA

LISTE STANDARD

3,7 anni



Adulto


3,4 anni



Pediatria


LISTE URGENZE NAZIONALI*

0,6 mesi



Adulto

5,5 mesi



Pediatria



656

Totale pazienti
in lista al
31/12/2021

Iscrizioni
al 31/12/2020


661

Ingressi 2021


379

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Opzioni chirurgiche per l'insufficienza cardiaca avanzata



Trapianto Cardiaco “gold standard” ma severamente limitato dalla carenza di donatori



Assistenza Ventricolare Meccanica

Come Ponte al Trapianto o Terapia Permanente

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

2019 EACTS Expert Consensus on long-term mechanical circulatory support

Evgenij V. Potapov^{a,*} (EACTS Chairperson), Christiaan Antonides^{b,†},
 Maria G. Crespo-Leiro^c, Alain Combes^{d,e}, Gloria Färber^f, Margaret M. Hannan^g, Marian Kukucka^h,
 Nicolaas de Jongeⁱ, Antonio Loforte^j, Lars H. Lund^k, Paul Mohacsi^l, Michiel Morshuis^m, Ivan Netukaⁿ,
 Mustafa Özbaran^o, Federico Pappalardo^p, Anna Mara Scandroglia^q,
 Martin Schweiger^r, Steven Tsui^s, Daniel Zimpfer^t and Finn Gustafsson^{u,*} (EACTS Chairperson),
 The Task Force on Long-Term Mechanical Circulatory Support of the EACTS



Recommendations for evaluation and selection of patients for LT-MCS therapy

Recommendation	Class	Level	References
It is recommended that reversible causes of heart failure are ruled out.	I	B	
LT-MCS implantation should be considered in patients with the following: • New York Heart Association functional class III–IV and • Ejection fraction $\leq 25\%$ and At least one of the following criteria: ◦ INTERMACS 2–4 ◦ Inotrope dependence ◦ Progressive end-organ dysfunction ◦ Peak $\text{VO}_2 < 12 \text{ ml/kg/min}$ ◦ Temporary MCS dependence	IIa	B	
LT-MCS implantation may be considered in patients with: • New York Heart Association functional class III–IV and • Ejection fraction $\leq 25\%$ and ◦ To reverse elevated pulmonary vascular resistance or potentially reversible renal failure in potential heart transplant candidates ◦ To allow time for transplant contraindications to be reversed such as recent cancer, obesity and recovering drug and alcohol dependence in potential heart transplant candidates	IIb	B	



LVAD evolution




**1° generation
“volume
displacement
pump”**



**2°
generation
“continous
flow rotary
pump”**






**3° generation
“FullMagnetic
levitation”**



HeartMate III LVAD

Full MagLev Technology

- Continuous-flow rotary technology
- Centrifugal configuration
- Non contact bearing design
- Development of pulsatile flow
- Better “washing” of the impeller with reduced risk of thrombus formation.
- Better surface biocompatibility



I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa



JAMA Cardiology | Original Investigation

Association of Clinical Outcomes With Left Ventricular Assist Device Use by Bridge to Transplant or Destination Therapy Intent

The Multicenter Study of MagLev Technology in Patients Undergoing Mechanical Circulatory Support Therapy With HeartMate 3 (MOMENTUM 3) Randomized Clinical Trial

Daniel J. Goldstein, MD; Yoshifumi Naka, MD, PhD; Douglas Horstmanhof, MD; Ashwin K. Ravichandran, MD, MPH; Jacob Schroder, MD; John Ransom, MD; Akinobu Itoh, MD; Nir Uriel, MD; Joseph C. Cleveland Jr, MD; Nirav Y. Rebecca Cogswell, MD; Erik E. Suarez, MD; Brian D. Lowes, MD, PhD; Gene Kim, MD; Pramod Bonde, MD; Farooq H. Sheikh, MD; Poornima Sood, MD; David J. Farrar, PhD; Mandeep R. Mehra, MD, MSc

Figure 3. Comparison of Hemocompatibility-Associated Adverse Events With HeartMate 3 vs HeartMate II Devices in the Bridge to Transplant (BTT)/Bridge to Transplant Candidacy (BTC) and Destination Therapy (DT) Cohorts



Survival at 2 years free of disabling stroke (defined as a modified Rankin score greater than 3) or reoperation to replace or remove a malfunctioning device in the bridge to transplant (BTT)/bridge to transplant candidacy (BTC) and destination therapy (DT) cohorts. HR indicates hazard ratio.


I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Primary results of long-term outcomes in the MOMENTUM 3 pivotal trial and continued access protocol study phase: a study of 2200 HeartMate 3 left ventricular assist device implants

Mandeep R. Mehra^{1*}, Joseph C. Cleveland Jr², Nir Uriel³, Jennifer A. Cowger⁴, Shelley Hall⁵, Douglas Horstmannshof⁶, Yoshifumi Naka³, Christopher T. Salerno⁷, Joyce Chuang⁸, Christopher Williams⁸, and Daniel J. Goldstein⁹, on behalf of the MOMENTUM 3 Investigators


¹Brigham and Women's Hospital Heart and Vascular Center and Harvard Medical School, Boston, MA, USA; ²University of Colorado School of Medicine, Aurora, CO, USA; ³Columbia University College of Physicians and Surgeons and New York-Presbyterian Hospital, New York, NY, USA; ⁴Henry Ford Hospitals, Detroit, MI, USA; ⁵Medical Center, Dallas, TX, USA; ⁶INTEGRIS Baptist Medical Center, Oklahoma City, OK, USA; ⁷St. Vincent Heart Center, Indianapolis, IN, USA; and ⁸Mo

European Journal of Heart Failure 2021



A Study of 2200 HeartMate 3 Left Ventricular Assist Device Implants in the MOMENTUM 3 Trial Portfolio

Are long-term outcomes with the HM3 LVAD in the post-pivotal trial experience different from early pivotal-trial observations?



Year Endpoints


Composite Endpoint
Survival free of disabling stroke or reoperation to replace or remove a malfunctioning device

Overall Survival

Overall Adverse Event Burden
Device pump thrombosis, stroke, bleeding, infection, right heart failure, cardiac arrhythmia, respiratory failure, and renal dysfunction

Cause Readmission Burden

Principal Outcomes



- 2200 HM3 implanted patients - 515 pivotal trial and 1685 continuous access protocol (CAP) patients
- Similar 2-year survival between CAP and pivotal trial cohorts (81.2% vs 79.0%) despite sicker patients (more intra-aortic balloon pump use and INTERMACS 1 profile) and more transplant ineligible patients intended for destination therapy in CAP
- Similar 2-year survival between CAP and pivotal trial cohorts in transplant ineligible patients (79.1% vs 76.7%) even after adjusting for baseline differences (HR=0.89 [95% CI: 0.68-1.16], P=0.38)

Accumulating post-pivotal trial experience with the HM3 LVAD suggests a lower adverse event burden, reduced hospitalizations and similar survival free of disabling stroke or reoperation to replace or remove a malfunctioning pump as compared to the pivotal MOMENTUM 3 trial outcomes at 2 years

These beneficial outcomes were noted across the continuum of clinical severity in advanced heart failure and especially among transplant ineligible patients in whom outcomes may now compare favorably with those in transplant eligible patients at 2 years


I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

JAMA | Original Investigation

Five-Year Outcomes in Patients With Fully Magnetically Levitated

Centrifugal-Flow Left Ventricular Assist Devices vs Axial-Flow Left Ventricular Assist Devices (LVADs)


Figure 2. Composite End Point and Overall Survival in a Study of 5-Year Outcomes in Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices (LVADs)



JAMA | Original Investigation

Five-Year Outcomes in Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices in the MOMENTUM 3 Randomized Trial

Figure 3. Serious Adverse Events in a Study of 5-Year Outcomes in Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices




JAMA | Original Investigation

Five-Year Outcomes in Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices in the MOMENTUM 3 Randomized Trial

Mandeep R. Mehra, MD, MSc; Daniel J. Goldstein, MD; Joseph C. Cleveland, MD; Jennifer A. Cowger, MD, MS; Shelley Hall, MD; Christopher T. Salerno, MD; Yoshifumi Naka, MD, PhD; Douglas Horstmannshof, MD; Joyce Chuang, PhD; AiJia Wang, MPH; Nir Uriel, MD, MSc


Figure 4. Cause-Specific Mortality 5 Years After Implant in a Study of Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices



I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

LVAD Survival Across Intermacs[†] Database Eras¹

REAL-WORLD OUTCOMES
FOR THE HEARTMATE 3[™]
LVAD ARE CONSISTENT
WITH MOMENTUM 3
OUTCOMES²



Data presented for informational purposes only.

1. Adapted from Teuteberg, et al. *Ann Thorac Surg*. 2020;109:649-60.

2. Mehra, et al. *N Engl J Med*. 2019;380:1618-1627.

3. Khush, et al. The International Thoracic Organ Transplant Registry of the International Society for Heart and Lung Transplantation: Thirty-fifth Adult Heart Transplant Report. 2018.

4. Kirklin, et al. *J Heart Lung Transplant*. 2014;33:555-564.

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Long-term mechanical circulatory support (destination therapy): On track to compete with heart transplantation?

James K. Kirklin, MD,^a David C. Naftel, PhD,^a Francis D. Pagani, MD, PhD,^b Robert L. Kormos, MD,^c Lynne Stevenson, MD,^d Marissa Miller, DVM, MPH,^e and James B. Young, MD^f


Objectives: Average 2-year survival after cardiac transplantation is approximately 80%. The evolution and subsequent approval of larger pulsatile and, more recently, continuous flow mechanical circulatory support (MCS) technology for destination therapy (DT) offers the potential for triage of some patients awaiting cardiac transplantation to DT.

Methods: The National Heart, Lung, and Blood Institute Interagency Registry for Mechanically Assisted Circulatory Support (INTERMACS) is a national multi-institutional study of long-term MCS. Between June 2006 and December 2011, 127 pulsatile and 1160 continuous flow pumps (24% of total primary left ventricular assist devices [LVADs]) carried an initial strategy of DT therapy.

Results: By multivariable analysis, risk factors ($P < .05$) for mortality after DT included older age, larger body mass index, history of cancer, history of cardiac surgery, INTERMACS level I (cardiogenic shock), dialysis, increased blood urea nitrogen, use of a pulsatile flow device, and use of a right ventricular assist device (RVAD). Among patients with a continuous flow LVAD who were not in cardiogenic shock, a particularly favorable survival was associated with no cancer, patients not in cardiogenic shock, and blood urea nitrogen less than 50 mg/dL, resulting in 1- and 2-year survivals of 88% and 80%. ←

Conclusions: (1) Evolution from pulsatile to continuous flow technology has dramatically improved 1- and 2-year survivals; (2) DT is not appropriate for patients with rapid hemodynamic deterioration or severe right ventricular failure; (3) important subsets of patients with continuous flow DT now enjoy survival that is competitive with heart transplantation out to about 2 years. (J Thorac Cardiovasc Surg 2012;144:584-603)


The Future of VAD vs Medical Therapy



Patient Profile/ Status: INTERMACS Levels

1. Critical cardiogenic shock
2. Progressive decline
3. Stable but inotrope dependent
4. Recurrent advanced HF
5. Exertion intolerant
6. Exertion limited
7. Advanced NYHA III

INDICATIONS for Transplant/ VAD



AMBULATORY HEART FAILURE PATIENTS

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Evaluation for Heart Transplantation and LVAD Implantation

JACC Council Perspectives

Maya Guglin, MD
Maryl R. Johnson, MD
Biykem Bozkurt, MD
and Leadership

ABSTRACT

Timely referral of patients with advanced heart failure is a key to good outcomes for advanced treatment decision making. The decision-making process for heart transplantation and LVAD implantation is a 3-step process. 1) Are there any contraindications to transplantation or LVAD? 2) Are there any medical conditions that would preclude transplantation or LVAD? 3) Are there any medical conditions that would preclude transplantation or LVAD?



CENTRAL ILLUSTRATION Advanced Heart Failure Center: Decision Making

TABLE 2 Simplified Signs Prompting Referrals to Advanced HF Center

I	Inotropes	Previous or ongoing requirement for dobutamine, milrinone, dopamine, or levosimendan
N	NYHA class/natriuretic peptides	Persisting NYHA functional class III/IV and/or high BNP or NT-proBNP
E	End-organ dysfunction	Worsening renal or liver dysfunction
E	Ejection fraction	Very low ejection fraction (<25%)
D	Defibrillator shocks	Recurrent appropriate defibrillator shocks
H	Hospitalizations	At least 1 hospitalization with HF in the past 12 months
E	Edema/escalating diuretic agents	Persistent fluid overload and/or increased diuretic requirement
L	Low BP	Consistently low BP (systolic <90 to 100 mm Hg)
P	Prognostic medications	Inability to up-titrate (or need to decrease/cease) ACE inhibitors, beta-blockers, ARNIs, or MRAs



- Timely referral is a key to good outcomes of transplantation and LVAD placement.

Eligible for LVAD only

Implant LVAD

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Evaluation for Heart Transplantation and LVAD Implantation



JAC

Maya
Maryl
Biyker
and Le

ABS

Time
patie
obsc
for a
mak
sync
3-st
2) A
assis
(J A

yes

Systemic illness
Liver cirrhosis
Malignancy with poor prognosis
Dementia or psychiatric illness
Financial restraints
Poor social situation

**Continue medical
management ± inotropes
or palliative therapy**

yes

osis
ess

s

Is Transplant/LVAD contraindicated?

Is Transplant/LVAD contraindicated?

no

Transplant or LVAD?

Eligible for transplant or LVAD

List for transplant but
implant LVAD if/when unstable

Eligible for transplant but not LVAD


List for transplant

Eligible for LVAD but not
transplant, may become
a transplant candidate in the future

Implant LVAD and
reconsider transplant later

Eligible for LVAD only


Implant LVAD




I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Pre-LVAD Frailty

Post-LVAD Frailty





Patient A



Favorable Outcome

Lower risk for premature death or complications, with marked improvement in functional status



Patient B



Intermediate Outcome

Moderate risk for premature death and complications with some persistent functional limitation

Patient C



Unfavorable Outcome

High risk for premature death and complications with failure to improve functional status

4 Percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Maria G. (

icolaas de

Martin :

Recommendation	Class	Level	References
Renal function			
In patients with renal dysfunction, optimization via improvement of cardiac output and reduction of filling pressures is recommended.	I	B	[56]
Liver function			
Liver function evaluation with bilirubin is recommended.	I	B	[57, 58]
In patients with increasingly elevated bilirubin levels, temporary MCS, ahead of possible LT-MCS implantation, may be considered.	IIb	B	[59]
Pulmonary function			
Treatment of preimplant pulmonary oedema is recommended before implantation.	I	B	[60, 61]
Left ventricular unloading on extracorporeal life support to optimize lung function should be considered.	IIa	B	[62]
Respiratory physiotherapy should be considered.	IIa	C	
Coagulation			
Withdrawal of dual antiplatelet therapy and/or vitamin K antagonists to reduce the risk of bleeding is recommended.	I	B	[63, 64]
The use of short-acting intravenous anticoagulation as bridging is recommended.	I	B	[64]
Administration of procoagulants shortly before implantation of the LT-MCS may be considered.	IIb	B	[64]
Optimization of coagulation prior to surgery should be considered, especially in patients on temporary MCS.	IIa	C	
Nutritional, metabolic and endocrine considerations			
Preoperative assessment of metabolic, endocrine and nutritional status, including possible interventions for arising issues, should be considered.	IIa	C	
Nutritional support, if necessary, may be considered.	IIb	C	[65, 66]

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Evaluation for Heart Transplantation and LVAD Implantation

JACC Council Perspectives

Maya Guglin, MD
Maryl R. Johnson, MD
Biykem Bozkurt, MD
and Leadership C

ABSTRACT


Timely referrals
patients with ac
obscured from r
for advanced th
making. The doc
syndrome of lov
3-step process a
2) Are there con

*Systemic illne
Liver cirrhosis
Malignancy v
Dementia or
Financial rest
Poor social si*

*Contir
managem
militi*

BRIDGE to TX

DESTINATION




Needed Advancements

- No driveline
- Wireless
- Improved hemocompatibility
- Ability to free disconnected

EvaHeart LVAD

- Hydraulically suspended impeller system promotes pulsatility
- More physiologically responsive
- COMPETENCE trial ongoing



CoreWave LVAD

- «wave membrane» technology
- Physiologic mimickig of the native heart allows for pulsatile flow
- Pre-clinical studies



When LVAD? When BiVAD/THA?

Key decision when assessing patients for VAD indication. Right ventricular failure during LVAD support is correlated with increased morbidity and mortality

Predictive index of right ventricular failure

- **Ecografici:**
 - TAPSE < 14 mm
 - RV ED S/L axis ratio > 0.57 (indice di sfericità)
 - FAC < 31%
 - strain della parete laterale (PSSrL < 0.6/s)
 - TR > II
 - RV/LV ratio > 0.75

Predictive index of right ventricular failure

- Emodinamici:
 - $CVP > 15 \text{ mmHg}$
 - $RAP/PCWP > 0.63$
 - $RVSWI < 300 \text{ mmHg/ml/m}^2 \rightarrow (MAP - RAP) \times SVI = 300-900 \text{ (v.n.)}$
 - $PAPi < 1.85 \rightarrow (PAS - PAD)/RAP$
 - $CI < 2.2$
 - $DP \text{ RV-RA} < 35 \text{ mmHg}$

Prediction of right ventricular assist device review and meta-analyses

Diego Bellavia^{1*}, Attilio Iannace¹, Michele Pilato¹, Sudhir S. Iyer¹, Valentina Agnese¹, Calogero M. D'Amico¹, Michael Dandel⁵

Table 4 Summary of the most relevant risk factors for right ventricular failure

Predictor	Comment
Demographic and clinical characteristics	
Age	Although several studies have reported that RVF patients are younger, according to meta-analysed data, age is a non-significant predictor of RVF.
Female gender	Females have apparently higher risk of developing post-LVAD RVF compared with males; however, this is specifically true in reports where primarily PF-LVADs are implanted.
Need for mechanical ventilation	Patients needing ventilatory support have roughly threefold increase in RVF risk, independently of type of implanted LVAD or gender. This predictor has good specificity but very low sensitivity.
Need for dialysis/CRRT	Patients needing dialysis have the highest risk of developing RVF after LVAD implantation, although this observation is based on relatively few studies ($n = 4$).
Biomarkers	
High INR	INR represents the single most significant and homogeneous biomarker in risk stratifying RVF in LVAD recipients, even a small increase should warrant attention and re-assessment.
High NT-proBNP level	Although NT-proBNP is higher in patients developing RVF, this observation is highly heterogeneous and overall marginally significant.
Haemodynamic parameters (by right heart catheterization)	
Low RVSWI	Role of RVSWI as a predictor of RVF after LVAD has been confirmed. However, its usefulness is significantly less important in the context of CF-LVADs compared with PF-LVADs.
High central venous pressure	Together with RVSWI, it is the haemodynamic predictor with the greatest effect size, specifically in patients undergoing CF-LVAD implantation.
Echocardiography	
Moderate-to-severe RVD	This relatively simple, pure qualitative assessment of RV performance is extremely predictive of RVF and should be part of routine echocardiographic evaluation in patients considered for LVAD.
High RV/LV ratio	To date, the RV/LV ratio obtained in a four-chamber apical view is the single most significant quantitative echo measurement able to define risk of RVF in LVAD recipients.
Low longitudinal sS of the RV free wall	Although sS of the right ventricle is very promising, and probably useful, there is too much variability in measures and there are too few published studies to reach a definitive conclusion.

CF-LVAD, continuous-flow left ventricular assist device; CRRT, continuous renal replacement therapy; INR, international normalized ratio; LV, left ventricular; LVAD, left ventricular assist device; NT-proBNP, N-terminal pro-brain natriuretic peptide; PF-LVAD, pulsatile flow-left ventricular assist device; RV, right ventricular; RVD, right ventricular dysfunction; RVF, right ventricular failure; RVSWI, right ventricular stroke work index; sS, systolic strain.

Prediction of right ventricular failure by echocardiographic measurements of right ventricular assist device review and meta-analysis studies

Diego Bellavia^{1*}, Attilio Iacovoni², C Michele Pilato¹, Sudhir S. Kushwaha³, Valentina Agnese¹, Calogero Falletta⁴, Michael Dandel⁵





Figure 8 Forest plot of the pooled estimate of the most significant echocardiographic measurements of right ventricular failure (RVF). (A) Forest plot of the pooled estimate of the impact of right ventricular/left ventricular basal diameter ratio (RV/LV ratio) on RVF (analysis not stratified). (B) Forest plot of the pooled estimate of the impact of RV free wall longitudinal systolic strain (sS) on RVF (analysis not stratified). CI, confidence interval; SMD, standardized mean difference.

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa






Rame, J.E. et al. J Am Coll Cardiol. 2021;78(23):2294-2308.

Patients from the Society of Thoracic Surgeons Intermacs database who survived to 3 months on left ventricular assist device (LVAD) support alone were divided into categories on the basis of right heart failure (RHF) status at 3 months. Compared with patients with mild and moderate right heart failure, those with no right heart failure had a significantly lower 12-month cumulative incidence of death, rehospitalization, stroke and gastrointestinal (GI) bleeding.






HeartMate3 Bi-VAD

Bridge to Transplantation






I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

CardioWest TAH



I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

CARMAT TAH



Biological contact surfaces →

No shear stress⁴

Fill-to-empty displacement pump

- 4 biological valves
- 2 hybrid membranes
 - Blood side: layer of bovine pericardium
 - Pump side: layer of polyurethane
- Cavities' walls covered by ePTFE


I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

FEATURED PAPERS

Interagency registry for mechanically assisted circulatory support report on the total artificial heart

J Heart Lung Transplant 2018;37:1304–1312

Intermecs Implants: June 2006 – December 2014




All Primary implants for left ventricular support
n= 13286

TAH N=301

Pulsatile Flow
(LVAD + RVAD)
N=339


Continuous
Flow
(LVAD + RVAD)
N=358



Pulsatile BIVAD including Total Artificial Hearts (TAHs) represented around 3% of the total devices implantation in 2014

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

La rete Scompenso Cardiaco Avanzato



I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Heart Failure Center con Heart Team

Multidisciplinare dedicato

Percorsi
decisionali

Condivisione
di strategie



Percorso decisionale
personalizzato e
risultato ottimale del
trattamento



Cardiologo
territorio



Team
Multidisciplinare



Cardiologo
Centro Spoke



Heart Failure
Team



Heart Failure
Center

Conclusioni

- Il **Trapianto** è il **Gold Standard** per il trattamento dello scompenso cardiaco refrattario.
- **L'Assistenza Ventricolare Meccanica** è oggi una soluzione terapeutica superiore alla terapia medica in molti pazienti con scompenso avanzato in attesa di Tx.
- Una collaborazione stretta tra cardiologo e **Heart Failure Center** è determinante per una gestione ottimale del paziente nel suo percorso clinico, per l'indicazione a trapianto o VAD e alla scelta del Timing chirurgico.

I percorsi diagnostico-terapeutici del paziente con insufficienza cardiaca avanzata: chi LVAD, chi trapianto, chi terapia palliativa

Grazie per l'attenzione



AZIENDA OSPEDALIERA
SAN CAMILLO FORLANINI

